

LAMDOT TONE

Comprehensive Liquid Mineral & Amino Acid Veterinary Feed Supplement

Description

This veterinary feed supplement is a carefully balanced nutrient formulation designed to provide essential minerals and amino acids to poultry. It supports optimal growth, bone development, metabolic functions, and immune health. The supplement combines key macro and micro minerals such as calcium, phosphorus, iron, copper, cobalt, zinc, manganese, sodium, potassium, sulphur, chloride, magnesium, iodine, alongside important amino acids and vitamins like lysine, biotin, and selenium. This synergistic blend promotes enhanced productivity and overall well-being in poultry flocks.

Specification

- Calcium: 4.5 gm
- Phosphorus: 2.5 gm
- Iron: 5 gm
- Copper: 40 gm
- Cobalt: 65 gm
- Zinc: 70 mg
- Manganese: 60 mg
- Sodium: 50 gm
- Potassium: 2.50 gm
- Sulphur: 4.50 gm
- Chloride: 1 mg
- Magnesium: 12.5 gm
- Iodine: 1 mg
- Lysine: 32.5 gm
- Biotin: 1 ppm (parts per million)
- Selenium: 0.5 mg

Category & Pack Size

- Category: Poultry
- Pack Size: 1 litre

Indication

This feed supplement is indicated for the nutritional support and prevention of mineral and amino acid deficiencies in poultry. It promotes healthy bone structure, improves metabolic processes, enhances immune response, and supports overall growth and production efficiency.

Dosage and Administration

- Dosage: Add the recommended quantity to the daily feed ration as per veterinary guidance; typically 1-2 ml per bird per day or as advised.
- Administration: Oral, mixed thoroughly in feed or drinking water.

Product Form

- Liquid feed supplement.

Usage

Recommended for all stages of poultry growth to maintain mineral balance, support metabolic health, and enhance production during stress, growth spurts, or recovery periods.

Advantages

- Complete mineral and amino acid profile for balanced nutrition.
- Improves growth rates and feed conversion efficiency.
- Boosts immune function and disease resistance.
- Easy to incorporate into existing feeding programs.

Adverse Effects

- Generally safe when used as directed.
- Overuse may cause mineral imbalances; avoid exceeding recommended dosage.
- Consult a veterinarian if adverse reactions occur.

Schedule

Non-controlled veterinary feed supplement; use as per veterinary advice.

Precautions

- Store in a cool, dry place away from direct sunlight.
- Shake well before use.
- Keep out of reach of children and unauthorized persons.
- Use clean equipment to avoid contamination.

Withdrawal Period

Follow local veterinary guidelines; typically a withdrawal period of 7-14 days before slaughter or egg collection to ensure safety.

Storage and Handling

- Store between 2°C and 25°C.
- Protect from freezing and heat.
- Seal tightly after use.

HS & HSN Codes

- HS Code: 2309.90 (Preparations of a kind used in animal feeding)
- HSN Code: 23099000

Special Instructions

- Use only under veterinary supervision.
- Maintain hygiene during handling and administration.
- Monitor birds for any unusual signs post-administration.

Details of Human and Veterinary Application

- Exclusively for veterinary use in poultry.
- Not intended for human consumption.
- Avoid contact with skin and eyes; wash hands after use.

? Frequently Asked Questions (FAQs)

Q1: What is the full form of ppm?

A1: Parts Per Million (ppm) is a unit measuring very small concentrations of substances.

Q2: Can this supplement be used for other animals?

A2: This product is formulated specifically for poultry; consult a veterinarian before use in other species.

Q3: How soon will benefits be seen?

A3: Noticeable improvements typically appear within 7-14 days depending on the bird's condition.

Q4: Can it be mixed in both feed and water?

A4: Yes, it can be safely mixed into feed or drinking water for ease of administration.

Q5: Is there a withdrawal period for eggs and meat?

A5: Yes, follow the recommended withdrawal period (usually 7-14 days) before eggs or meat are used for consumption.

